

## **+ IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW:**

COPIKTRA can cause fatal and/or serious toxicities including treatment-related mortality, infections, diarrhea or colitis, cutaneous reactions, and pneumonitis. Contact the healthcare provider on this card **immediately** if your patient reports any signs or symptoms.

## **FOR HEALTHCARE PROVIDERS**

 I am taking COPIKTRA for my treatment of leukemia or lymphoma.

## **+ IMPORTANT SAFETY INFORMATION FOR PATIENTS TAKING COPIKTRA®**

\_\_\_\_\_  
Patient Name

\_\_\_\_\_  
COPIKTRA Prescriber Name

\_\_\_\_\_  
Prescriber Phone Number



**PATIENT SAFETY  
WALLET CARD**



**Show this to any doctor  
involved in your care**

## **+ IMPORTANT SAFETY INFORMATION**

### **FOR THE PATIENT**

COPIKTRA can cause death due to treatment side effects. The serious side effects that may lead to death include:

- Infections
- Diarrhea or inflammation of your large intestine (colitis)
- Skin reactions
- Inflammation of the lungs

If you have any of these symptoms, call your healthcare provider **right away**:

- Fever, chills, or other signs of infection
- New or worsening diarrhea
- Stool with mucus or blood
- Severe stomach-area (abdominal) pain
- New or worsening skin rash or other skin reactions, including:
  - Painful sores or ulcers on your skin, lips, or in your mouth
  - Severe rash with blisters or peeling skin
  - Rash with itching
  - Rash with fever
- New or worsening cough or difficulty breathing



**IMPORTANT TO REMEMBER:** Call your doctor or get emergency medical care right away if you have any of these symptoms! Show this card to any doctor involved in your care!